

Scientific Combination of DNA Evidence: A Handgun Mixture in Eight Parts

Mark W Perlin, PhD, MD, PhD
Cybergenetics, Pittsburgh, PA, USA

Matt Greenhalgh
Orchid Cellmark, Abingdon, UK

Australia and New Zealand
Forensic Science Society
September, 2010




Cybergenetics © 2003-2010


Handgun Evidence







DNA Swabs












Computer Infers Major Genotype



Definite genotype
at every locus
100% probability
at one allele pair

Full RMP
 $\log(LR) = 16.72$
trillion-fold increase
 $4.23 \rightarrow 16.72$
over CPI

Computer Infers Minor Genotype




Uncertain genotype
at every locus
< 100% probability
at multiple allele pairs

Finds match strength
 $\log(LR) = 4.99$
100,000-fold increase
 $0 \rightarrow 4.99$
over no result


Joint Likelihood Function

Locus D18

85% major [15 19]
15% minor [13 13]




Infer Minor Genotype




More genotype certainty with
two amplifications of
one PCR template

Greater match strength
 $\log(LR) = 5.25$
some increase
0 → 4.99 → 5.25

Mixture Weight: 4 templates



Joint Likelihood Function



Infer Minor Genotype




Higher genotype certainty with
two amplifications of
two PCR templates

Greater match strength
 $\log(LR) = 9.73$
large increase
0 → 4.99 → 5.25 → 9.73

Joint Likelihood Function


- 3
- 4
- 5
- 6



Locus D18




major [15 19]
minor [13 13]


Infer Minor Genotype



Definite genotype at every locus
two amplifications of
four PCR templates

Full match strength
 $\log(LR) = 12.39$
reaches RMP
0 → 4.99 → 5.25 → 9.73 → 12.39





Observations

- Coancestry - half a log unit
- Three unknown contributors
- Multiple amplifications are good, multiple templates are better
- Data constrains genotypes; more data, more constraints
- *Complementarity*: combining "unlike data" is more informative

Conclusions

Two mixture genotype examples

Major contributor Minor contributor	Combining DNA evidence using a joint likelihood function is <i>more informative</i> than separately examining data in isolation
--	--

Trillion-fold increase in identification information
